

University of Groningen

A Closer Look at the Anatomy of Spiral Galaxies

Peters, Stephan

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2014

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Peters, S. (2014). *A Closer Look at the Anatomy of Spiral Galaxies*. [Thesis fully internal (DIV), University of Groningen]. [S.n.].

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

- The near constant value of the maximum surface brightness of the HI along a significant fraction of the major axis in many edge-on galaxies is a clear indication of HI self-absorption. (Chapters 1 and 2)
- Treating the HI of an edge-on galaxy as optically thin will underestimate the HI mass of a galaxy by at least 23 to 30%. If the HI is fragmented into clouds, then even more HI can remain hidden. (Chapter 2)
- The assumption that the HI is optically thin or self-absorbing has a profound impact on the measured scale-height, face-on surface density and velocity dispersion of the gas. (Chapter 3)
- It is very hard to measure the velocity dispersion of the HI in edge-on galaxy accurately (Chapter 3).
- The measured shape of the dark matter halo through observations of edge-on galaxies is very susceptible to errors due to the many underlying assumptions. (Chapter 5)
- Given the assumptions and uncertainties involved in working with observations, one should always use advanced optimization techniques that allow these errors to be estimated.
- Scattered light from the stellar disk of a face-on galaxy will often look like a stellar halo around that galaxy. This light can hide the signature of a truncation in the stellar disk. (Chapter 7)
- Science is in need of a very thorough revision of its employment and career strategy.
- As astronomy transitions ever more into a desk job, we run the risk of forgetting the feeling of magic and wonder one gets from simply looking at the night sky.
- Ink is the new black gold.